

2 февраля 2009 г.

Optical Identifications of 230 HIPASS Radio Sources

I. D. Karachentsev,¹ D. I. Makarov,^{1,2} V. E. Karachentseva,³ and O. V. Melnik³

¹*Special Astrophysical Observatory, Russian Academy of Sciences,
Nizhnii Arkhyz, 357169 Karachaevo-Cherkesskaya Republic, Russia**

²*Invited researcher in Lyon Observatory, Lyon, France*

³*Astronomical Observatory, Kiev National University,
Observatorna vul. 3, Kiev, 04053 Ukraine*

We present the coordinates, apparent magnitudes, and morphological types for 230 galaxies presumably identified with HIPASS (HI Parkes All-Sky Survey) sources. The new optical counterparts of the HIPASS sources follow the well-known statistical relationships between the hydrogen mass, luminosity, and type of galaxies. Low-surface-brightness galaxies constitute a significant fraction among these objects. The median value of the hydrogen mass-to-luminosity ratio for them is a factor of 2 or 3 higher than that for bright HIPASS galaxies, reaching $1.7M_{\odot}/L_{\odot}$. A number of our objects are located near the boundary $\log(M_{HI}/L_B) = 0.2(M_B + 20)$ that defines the zone of gravitational stability of disk galaxies against large-scale star formation.

Key words: dwarf galaxies, galaxy evolution, interstellar gas in galaxies.

I. INTRODUCTION

The blind survey of the southern sky in the 21-cm HI line carried out with the 64-m radio telescope in Parkes (Australia) has revealed 4315 radio sources. The catalogs of HIPASS sources were published by Meyer et al. (2004), Koribalski et al. (2004), and Henning et al. (2000). The effective depth of this survey in radial velocities is $\sim 3000 \text{ km s}^{-1}$, i.e., most of the HIPASS sources are located in the Local Supercluster.

Numerous optical identifications of HIPASS objects were made by Staveley-Smith et

*Electronic address: ikar@sao.ru

al. (1998), Ryan-Weber et al. (2002), and Doyle et al. (2005). The latest list of optical identifications (<http://hipass.aus-vo.org>) contains probable optical counterparts (galaxies) for 3618 HIPASS sources, i.e., 84% of their total number. The remaining 16% of the objects have no optical identifications for various reasons: (a) two or more galaxies fall into the wide ($\sim 15'$) beam of the radio telescope; (b) objects are located at low Galactic latitudes in the zone of strong absorption; (c) a HIPASS source is a dwarf galaxy rich in gas but with a very low surface brightness similar to those detected by Karachentseva and Karanchentsev (1998), KK, Karachentseva and Karanchentsev (2000), KKs, and Karachentsev et al. (2000), KKSG, during an all-sky survey on POSS-II images is a HIPASS source. Recently, Wong et al. (2006) published a northern extension of the Parkes Survey to a declination of $+25.5^\circ$, bringing the total number of HIPASS radio sources to 5317. For some of the northern HIPASS sources, the NASA Extragalactic Database (NED) gives identifications without any reference to their origin.

When preparing the list of galaxies within the Local Supercluster (with radial velocities up to 3800 km s^{-1}), we paid attention to the unidentified HIPASS sources. To supplement our list, we made an independent optical check for more than 500 HIPASS sources, which added over 200 more identifications. In this paper, we present data for 230 identified HIPASS sources, leaving without comments the cases where no optical counterparts were found.

II. RESULTS

When examining the Digitized Sky Survey areas with unidentified HIPASS sources, we took into account the intensity and width of the 21-cm line as well as the position of a particular galaxy relative to the position of the HIPASS object. We excluded the cases of a possible identification with several suitable optical galaxies. For each galaxy, we determined its morphological type T by taking into account its B and R brightness and visibility in the 2MASS survey as well as its total apparent B magnitude from comparison with the images of other galaxies of a similar structure with known magnitudes. Occasionally, we included galaxies with previously made identifications in our list by adding our own estimates of the morphological type and magnitude. Our identifications of 230 HIPASS sources are given in

the table¹. The columns of the table contain the following data: (1) HIPASS source name; in several cases, we added objects from the HIZOA (Donley et al. 2005), HIDEEP (Minchin et al. 2003), and HIJASS (Lang et al. 2003) surveys; (2) coordinates of the optical galaxy for the epoch 2000.0; (3) heliocentric radial velocities from the 21-cm line in km s^{-1} ; (4, 5) full width at half maximum (FWHM) of the 21-cm line, in km s^{-1} , and flux F in the 21 cm line, in Jansky km s^{-1} , taken mainly from HOPCAT (Doyle et al. 2005); (6) total apparent magnitude from NED (given with an accuracy of 0.01^m) or from our estimates with an accuracy of about 0.5^m ; (7) galactic extinction in the B band from Schlegel et al. (1998); (8) hydrogen mass-to-luminosity ratio in units of the solar mass and luminosity, which does not depend on the galaxy distance; (9) morphological type in the de Vaucouleurs numerical scale; (10) notes with a description of structural features. In the notes, we also give the galaxy names from NED (from the catalogs and lists); in the references to the 2dF, 6dF, APM, 2MASS, and other surveys, the galaxy names are given without any coordinate part.

Figure 1 presents negative DSS images of the four galaxies that we identified with HIPASS sources. The galaxies with a low surface brightness are at the center of each image; the image size is 6×6 arcmin.

III. DISCUSSION

Late-type and low-surface-brightness galaxies prevail among our identifications as “residual” with respect to the previous ones. About a quarter of the galaxies are located at low Galactic latitudes with extinction $A_b > 1.0^m$. Figure 2 shows the distribution of our identified galaxies in hydrogen mass-to-luminosity ratio on a logarithmic scale. The M_{HI}/L distribution has a median of $1.7M_\odot/L_\odot$; half of the entire sample is contained between the quartile values $(0.9 \div 2.4)M_\odot/L_\odot$. For comparison, note that the sample of 752 HIPASS BGC galaxies (Koribalski et al. 2004) has a median M_{HI}/L that is two and a half times smaller than that for the galaxies from the table. In Fig. 3, the M_{HI}/L ratio is plotted against the morphological type of the galaxies identified in our survey. As we see, more than half of the galaxies belong to the latest types: 10 (Ir), 9 (Im, BCD), and 7, 8 (Sd, Sdm). An expected positive correlation with the coefficient $R = 0.40$ and standard deviation 0.34 is

¹ The table is published completely only in electronic form and is accessible via <ftp://cdsarc.u-strasbg.fr/pub/cats/J> (130.79.128.5) or <http://cdsweb.u-strasbg.fr/pub/cats/J>.

seen between $\lg(M_{HI}/L)$ and T .

In Fig. 4, the hydrogen mass-to-luminosity ratio is plotted against the observed hydrogen line FWHM W_{50} (Wong et al. 2006) (uncorrected for the slope). As we see from the figure, the sources that we identified confirm the well-known fact of a decrease in the hydrogen mass fraction from faint (with smaller W_{50}) to bright galaxies. Note that all galaxies with $M_{HI}/L > 5$ lie in the range $W_{50} < 150 \text{ km s}^{-1}$.

Another expected relationship between $\lg(M_{HI}/L)$ and the absolute magnitude of a galaxy MB calculated from its radial velocity with $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is shown in Fig. 5. The solid line corresponds to the regression

$$\lg(M_{HI}/L) = 0.15(M_B + 17.5), \quad (1)$$

which reflects the well-known statistical relationships between the hydrogen mass of a galaxy and its radius, $M_{HI} \propto R^2$, and between the total luminosity and radius, $L \propto R^3$. Hence follows the relationship $M_{HI}/L \propto L^{-1/3}$, which is almost equivalent to (1). As we see from Fig. 5, there is no galaxy with a M_{HI}/L ratio smaller than $3 \cdot M_{\odot}/L_{\odot}$ among the dwarf galaxies fainter than B_T . Of course, the low accuracy of the BT estimate from DSS images introduces an additional scatter on the diagrams.

Warren et al. (2007) also constructed the relationship $M_{HI} \propto M_B$ for HIPASS galaxies and discussed it in connection with the problem of the minimum number of stars that can be formed in a dwarf system if it is stable. Considering HIPASS galaxies and invoking four known objects with detailed photometry and record high values of $M_{HI}/L > 5M_{\odot}/L_{\odot}$, (ESO 215-09, DDO 154, UGCA 292, NGC 3741), these authors found a positive correlation between M_{HI}/L and the luminosity of the galaxies that is almost identical to ours. Warren et al. (2007) pointed out the presence of an upper limit on this diagram,

$$\lg(M_{HI}/L)_{\max} = 0.20(M_B + 20), \quad (2)$$

under which the entire set of galaxies known to date lies. The authors explain the presence of this empirical limit as follows. According to Taylor and Webster (2005), the galactic gas disk in the absence of an internal stellar radiation field becomes gravothermally unstable, which gives rise to molecular hydrogen even for galaxies with very low baryon masses. The fraction of the unstable gas decreases from normal to progressively less massive galaxies. As a result, dwarf galaxies become stable at a smaller fraction of the gas transformed into stars.

Thus, the presence of the upper limit (2) can be interpreted as the existence of a minimum fraction of the baryon mass in a galaxy that must be transformed into stars for the galaxy to remain stable against large-scale star formation.

We can add four more galaxies from the Catalog of Nearby Galaxies (Karachentsev et al. 2004) with high values of (M_{HI}/L) to the four mentioned galaxies with record high ratios M_{HI}/L (ESO 215–09 (22), DDO 154 (9.4), UGCA 292 (7.0), and NGC 3741 (5.8)): HolmIX (11.0), ESO 564–030 (6.3), PGC 51659 (6.0), and AndIV (21.0). The hydrogen data for AndIV were taken from Chengalur et al. (2007). The hydrogen cloud in the Virgo Cluster (HI 1225+01) discovered by Giovannelli and Haynes (1989) with signatures of recent star formation (Salzer 1992) is also plotted on the diagram of Fig. 5. Based on the results of our work, we found ten more galaxies with a high gas content per unit luminosity: J0718–09 (6.6), J0727+04 (7.4), J1007–66 (12.4), Leib (19.1), KKSG 47 (7.8), J1549+16 (11.0), J1739–51 (10.4), J2011–09 (5.4), J2015+12 (6.6), and ESO 403–036 (5.1). Among them, only one galaxy, J1739–51, exceeds significantly the critical condition (2). However, its visual B_T estimate needs a photometric improvement.

Note that our sample includes the unusual dwarf system HIPASS J1123+13 in the nearby group around NGC 3628. We identified this HIPASS source with an object of an extremely low surface brightness with angular sizes of $1.4' \times 0.8'$ and a total apparent magnitude $B_T = 17.5^m$. According to Haynes et al. (1979), an HI filament, which is also barely seen optically, extends from this source to the neighboring bright galaxy NGC 3628. Obviously, this exotic object, which we called “Leib”, deserves a more detailed study at various wavelengths.

IV. ACKNOWLEDGMENTS

We wish to thank B. Koribalski and L.N. Makarova for advice and help in the work. This work was supported in part by the Russian Foundation for Basic Research (project nos. 07–02–00005, 08–02–00627, and 06–02–04017).

-
- [1] J. Chengalur, A. Begum, and I. D. Karachentsev, arXiv0711.2153 (2007).
 - [2] M. T. Doyle, M. J. Drinkwater, D. J. Rohde, et al., Mon. Not. R. Astron. Soc. 361, 34 (2005).
 - [3] J. L. Donley, L. Staveley-Smith, R. C. Kraan-Korteweg, et al., Astron. J. 129, 220 (2005).

- [4] R. Giovanelli and M. Haynes, *Astrophys. J.* 346, L5 (1989).
- [5] M. P. Haynes, R. Giovanelli, and M. S. Roberts, *Astrophys. J.* 229, 83 (1979).
- [6] P. A. Henning, L. Staveley-Smith, R. D. Ekers, et al., *Astron. J.* 119, 2686 (2000).
- [7] V. E. Karachentseva and I. D. Karachentsev, *Astron. Astrophys. Suppl. Ser.* 127, 409 (1998).
- [8] V. E. Karachentseva and I. D. Karachentsev, *Astron. Astrophys. Suppl. Ser.* 146, 359 (2000).
- [9] I. D. Karachentsev, V. E. Karachentseva, A. A. Suchkov, and E. K. Grebel, *Astron. Astrophys. Suppl. Ser.* 145, 415 (2000).
- [10] I. D. Karachentsev, V. E. Karachentseva, W. K. Huchtmeier, and D. I. Makarov, *Astron. J.* 127, 2031 (2004).
- [11] B. S. Koribalski, L. Staveley-Smith, V. A. Kilborn, et al., *Astron. J.* 128, 16 (2004).
- [12] R. H. Lang, P. J. Boyce, V. A. Kilborn, et al., *Mon. Not. R. Astron. Soc.* 342, 738 (2003).
- [13] M. J. Meyer, M. A. Zwaan, R. L. Webster, et al., *Mon. Not. R. Astron. Soc.* 350, 1195 (2004).
- [14] R. F. Minchin, M. J. Disney, P. J. Boyce, et al., *Mon. Not. R. Astron. Soc.* 346, 787 (2003).
- [15] E. Ryan-Weber, B. S. Koribalski, L. Staveley-Smith, et al., *Astron. J.* 124, 1954 (2002).
- [16] J. Salzer, *Astron. J.* 103, 385 (1992).
- [17] D. J. Schlegel, D. P. Finkbeiner, and M. Davis, *Astrophys. J.* 500, 525 (1998).
- [18] L. Staveley-Smith, S. Juraszek, B. S. Koribalski, et al., *Astron. J.* 116, 2717 (1998).
- [19] E. N. Taylor and R. L. Webster, *Astrophys. J.* 634, 1067 (2005).
- [20] B. E. Warren, J. Helmut and B. S. Koribalski, *Astron. J.* 134, 1849 (2007).
- [21] O. I. Wong, E. V. Ryan-Weber, D. A. Garcia-Appadoo, et al., *Mon. Not. R. Astron. Soc.* 371, 1855 (2006).

Таблица I: New probable optical identifications of HIPASS sources in the Local Supercluster (N = 230).

HIPASS (NED)	Optical counterpart	V_h	W_{50}	flux	B_t	A_b	M_{HI}/L	T	Comments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J0002−07	J000202.3−073913	3765	45	2.2	17.21	0.13	2.27	8	6dF
HIPASS J0002−15	J000225.8−155327	3416	94	3.7	16.7	0.12	2.40	6	
HIPASS J0021+08	J002041.7+083656	693	41	1.7	16.8	0.55	0.82	10	
HIPASS J0030−52	J003028.6−521515	2640	151	2.6	16.63	0.06	1.67	2	
HIPASS J0041−01b	J004139.7−020042	1949	76	3.7	16.89	0.12	2.86	10	
HIPASS J0116−63	J011650.0−633000	2306	130	2.5	16.7	0.09	1.66	7	
HIPASS J0207−06	J020724.4−061348	3819	82	3.4	17.2	0.11	3.53	10	
HIPASS J0227−15	J022720.8−152517	3767	254	10.3	14.85	0.09	1.25	7	MCG-03-07-19
HIPASS J0237+12	J023718.6+123108	960	51	1.9	16.4	0.63	0.59	9	
HIPASS J0258−14	J025812.6−142302	3272	152	6.4	16.47	0.18	3.19	8	APMUKS
HIPASS J0305−69	J030525.2−685614	1369	120	7.3	16.0	0.15	2.42	9	bluish, IRAS
HIPASS J0341+24	J034111.9+240046	1260	110	11.9	15.8	1.68	0.80	6	
HIPASS J0341+18	J034201.9+180831	1297	45	3.0	16.4	0.91	0.71	8	
HIPASS J0358+10	J035824.2+095845	1978	141	6.9	17.0	0.71	3.44	8	
HIPASS J0403−01	J040331.3−015540	937	92	9.5	16.4	0.93	2.21	10	LSB in a star halo
HIPASS J0419−21	J042012.7−211436	900	46	2.7	16.3	0.12	1.21	8	ESO 550-23
HIPASS J0421−51	J042036.0−511301	1046	142	5.9	16.5	0.06	3.37	9	LSBG F202-83?, 12'away
HIPASS J0423−56	J042251.7−561332	1356	49	1.1	16.25	0.08	0.49	10	APM
HIPASS J0438+00	J043846.1+001845	3302	51	2.2	17.1	0.19	1.94	10	
HIPASS J0458−07	J045839.7−073306	4052	87	3.7	17.0	0.27	2.75	8	
HIPASS J0501−45	J050103.8−451450	1140	76	2.5	16.06	0.08	0.93	8	APM
HIPASS J0519−04	J051915.6−042608	3848	53	3.1	16.8	1.12	0.88	8	
HIPASS J0526−61	J052617.8−612141	1219	101	7.7	15.3	0.21	1.27	8	
HIJASS J0529+72	J052920.5+722727	1089	46	11.0	16.0	0.44	2.81	10	HS98
HIPASS J0531+08	J053102.4+082104	961	90	5.4	16.9	1.89	0.82	10	
HIPASS J0544−16	J054423.6−162652	2158	152	3.9	16.1	0.38	1.15	7	
HIPASS J0554−71	J055441.2−715539	1482	64	3.1	16.9	0.32	2.01	9	
HIPASS J0559−42	J055954.3−425314	1013	106	5.2	16.4	0.26	2.25	10	

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J0605−02	J060552.2−022216	2462	126	5.2	17.9	1.6	2.61	10	
HIPASS J0619−52	J061925.9−524638	1116	59	4.9	15.2	0.34	0.65	10	
HIPASS J0620−27	J062007.1−272251	1789	47	3.0	16.8	0.17	2.03	10	
HIPASS J0620+20	J062103.5+201017	1318	138	8.7	18.0	3.47	0.86	7	ADBS (NED)
HIPASS J0626+24	J062621.0+243920	1473	99	8.3	17.6	1.79	2.65	7	
HIPASS J0630+23	J062958.2+233428	1452	135	11.0	17.1	1.18	3.87	7	
HIPASS J0635+11	J063548.5+111507	3575	77	4.0	17.5	4.36	0.11	7	
HIPASS J0643−25	J064350.2−254509	2505	111	4.0	16.5	0.43	1.63	6	
HIPASS J0646−23	J064600.5−230144	1896	48	2.7	18.0	0.52	4.02	10	
HIPASS J0647−15	J064732.4−151611	2792	132	3.5	17.5	2.59	0.49	7	
HIPASS J0650+16	J065036.3+162121	2549	80	5.3	16.5	0.64	1.77	8	
HIPASS J0653−73	J065352.7−734228	1208	94	3.5	17.0	0.58	1.96	7	
HIPASS J0700−02	J070005.3−022352	1774	52	3.1	17.8	3.72	0.20	10	
HIPASS J0702−28	J070220.0−281859	1801	82	3.9	17.0	0.51	2.33	10	[DDR2005]
HIZOA J0702+03	J070250.7+031114	3551	103	4.2	18.2	1.52	2.99	8	
HIZOA J0705+02	J070538.7+023720	1745	46	4.6	18.3	2.00	2.31	8	[H92]16ID
HIPASS J0712−39	J071228.9−390549	2566	60	3.4	16.8	0.99	1.09	8	
HIPASS J0712−41	J071310.6−411935	1689	48	2.5	16.9	0.72	1.12	8	
HIPASS J0713−30	J071331.7−303537	1740	74	3.0	17.0	0.61	1.63	7	
HIPASS J0716−40b	J071640.1−403305	3234	79	4.5	17.4	1.07	2.31	8	
HIPASS J0717−08	J071743.2−085517	2477	165	9.5	17.2	2.09	1.58	5	
HIPASS J0718−79	J071745.0−792440	1571	66	4.1	17.4	1.49	1.43	10	
HIPASS J0718−09	J071821.0−090316	909	57	13.2	18.2	1.90	6.61	8	
HIPASS J0721−30	J072106.2−300539	1951	105	6.6	17.0	0.90	2.75	7	
HIPASS J0725−24b	J072520.6−245643	2769	232	12.1	18.0	4.59	0.42	6	
HIPASS J0727+04	J072754.0+044146	2087	121	4.5	17.9	0.30	7.45	7	
HIPASS J0737−31	J073717.1−310654	2237	157	7.7	17.6	2.46	1.33	6	
HIPASS J0741−09	J074134.4−094201	2435	96	2.7	17.0	0.76	1.28	6	
HIPASS J0742−26	J074158.5−264248	2924	237	8.1	17.7	4.00	0.37	6	
HIPASS J0745−74	J074521.0−744820	1741	90	1.8	18.0	2.01	0.68	10	

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J0747–26b	J074712.8–261746	2580	170	19.0	17.1	2.54	1.92	6	
HIPASS J0751–37	J075136.9–371350	2808	87	10.3	18.2	5.26	0.23	7	
HIPASS J0752–49	J075218.7–493221	1158	56	2.6	16.7	0.89	0.83	8	
HIPASS J0754–38	J075448.4–385524	2757	270	16.4	17.2	3.47	0.77	7	
HIPASS J0758+10	J075812.3+110114	2346	94	5.7	16.4	0.10	2.86	7	
HIPASS J0759–71	J075923.5–715325	1368	110	2.4	17.5	1.11	1.31	10	
HIPASS J0801–21	J080125.1–215956	729	56	3.6	17.4	0.96	2.05	8	
HIZSS 035	J080953.9–414131	1995	318	24.0	16.9	5.23	0.17	6	
HIPASS J0834–18	J083414.2–184215	4161	147	6.7	15.11	0.34	0.82	4	[DDR2005]
HIPASS J0835–58	J083513.4–583646	1050	61	4.0	16.0	0.66	0.83	9	ESO 125-01
HIPASS J0836+05	J083633.8+051040	1866	60	2.6	17.5	0.12	3.53	10	
HIPASS J0838–61	J083827.1–612330	2973	91	5.9	16.5	0.83	1.66	7	
HIPASS J0844–33	J084355.0–330855	2316	201	10.6	16.8	1.65	1.84	7	
HIPASS J0846–76	J084624.1–765934	1520	50	2.4	16.9	0.67	1.13	10	
HIPASS J0848–33	J084903.4–330255	1320	116	4.2	17.8	1.4	2.31	7	
HIPASS J0848–26	J084906.0–261918	809	61	3.4	15.0	0.41	0.35	5	ESO 496-010
HIPASS J0854–54	J085432.4–544645	2729	59	3.8	17.1	1.51	0.99	8	
HIPASS J0856–11	J085651.8–110827	1879	68	3.0	17.0	0.22	2.33	9	
HIPASS J0857–29	J085705.6–290723	1967	33	3.9	17.1	0.78	2.00	8	
HIZSS 049	J085809.4–390953	1203	49	3.1	17.8	2.61	0.56	7	
HIPASS J0900–37	J090056.4–370723	1105	97	9.1	17.8	2.52	1.79	8	
HIPASS J0906–34	J090656.8–341910	1410	81	3.4	17.4	1.49	1.19	10	
HIPASS J0916–23b	J091658.0–231647	837	40	1.1	15.83	0.47	0.23	8	ESO 497-035
HIPASS J0917–34	J091741.7–344644	2366	154	8.4	17.4	1.19	3.87	8	
HIPASS J0920–34	J092044.6–342628	2410	157	7.0	14.9	1.10	0.35	5	ESO 372-015
HIPASS J0921–32	J092203.3–324717	1094	97	6.9	17.2	0.80	3.80	9	
HIPASS J0926–32	J092650.1–324748	1188	131	10.6	16.4	0.59	3.37	7	ESO 373-01
HIPASS J0939+00	J093857.1+004133	2076	66	3.3	16.5	0.30	1.51	10	SDSS
HIPASS J0941–15	J094059.4–153421	3619	154	4.9	16.2	0.25	1.79	6	
HIPASS J0941–02	J094101.8–024303	1633	128	4.5	16.9	0.15	3.40	8	

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J0942+04	J094250.9+045324	1955	66	2.9	17.0	0.24	2.21	8	
HIPASS J0944−31a	J094431.8−313320	2123	73	3.9	16.52	0.45	1.58	9	ESO 434-030
HIZSS 056	J094524.5−480828	882	79	7.9	17.0	2.09	1.10	8	
HIPASS J0946−15	J094622.7−154523	3164	82	3.5	16.5	0.28	1.63	10	
HIPASS J0952−49	J095310.9−491950	2258	100	4.0	17.0	2.08	0.56	7	
HIPASS J0958−29	J095813.6−292451	2311	127	14.0	15.9	0.36	3.47	5	6dF
HIPASS J0958−85	J100014.0−854140	1965	64	2.1	16.9	0.78	0.90	8	
HIPASS J1000−77	J100108.0−771826	1716	100	2.2	18.0	2.42	0.57	10	
HIPASS J1002−33	J100149.0−331542	2351	98	4.0	15.67	0.40	0.78	8	ESO 374-021
HIPASS J1003−49	J100300.7−492904	2411	107	7.5	17.3	2.26	1.18	7	
HIPASS J1005−28	J100539.5−282644	1037	100	9.0	15.74	0.34	1.98	7	ESO 435-039
HIPASS J1006−49	J100616.5−492829	2411	140	9.2	18.0	1.83	4.09	7	
HIPASS J1007−66	J100734.7−660635	1856	77	5.0	19.0	0.97	12.36	10	
HIPASS J1008−33	J100829.3−330835	1645	58	3.1	17.0	0.36	2.13	10	
HIPASS J1015−34	J101553.8−340653	2608	27	5.4	16.17	0.38	1.69	9	ESO 375-03
HIPASS J1016−39	J101607.2−395923	3006	69	8.1	16.6	0.57	3.16	8	
HIPASS J1024−12	J102428.3−122557	628	45	2.6	17.9	0.30	4.33	10	KKSG 19
HIPASS J1057−21	J105709.9−213612	2075	64	4.3	17.3	0.23	4.37	8	
HIPASS J1100−16	J110042.5−165436	3890	59	3.8	16.2	0.23	1.41	6	
HIPASS J1117−05	J111732.9−051639	3990	143	4.3	16.12	0.16	1.57	6	2dF
HIPASS J1122+13	J112313.5+134254	896	27	14.1	17.5	0.12	19.05	10	ELSB on POSS-IIB
HIPASS J1126−76	J112622.6−770012	2323	159	8.6	16.5	1.52	1.28	5	2MASS
HIPASS J1135−46	J113532.2−464310	2827	176	6.2	16.5	0.49	2.38	5	2MASS
HIPASS J1152−14	J115205.1−145330	2189	105	4.4	16.5	0.16	2.29	10	
HIPASS J1154+12	J115412.7+122604	1005	48	2.0	17.9	0.13	3.87	10	
HIPASS J1158−49	J115901.1−495102	3430	84	2.1	15.17	0.53	0.23	7	[DDR2005]
HIPASS J1201−15	J120141.0−150630	1364	77	4.4	15.0	0.21	0.55	9	MCG-02-31-011
HIPASS J1204−22	J120412.8−220635	1695	130	5.1	16.39	0.22	2.27	9	NPM1G-21.0322
HIPASS J1215+12	J121503.9+130155	2090	25	2.1	16.4	0.14	1.02	8	VCC0132
HIPASS J1215−17	J121545.1−173806	1391	102	5.1	16.0	0.27	1.51	10	in a star halo

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J1221−36	J122124.5−360638	2598	47	3.2	16.8	0.35	1.84	7	
HIPASS J1225−42	J122501.6−430245	2138	59	3.7	14.5	0.40	0.24	5	IRAS
HIPASS J1227−50	J122706.8−502031	3557	74	2.2	15.97	0.89	0.36	7	[DDR2005]
HIPASS J1228−50	J122858.4−500916	3694	149	5.8	14.9	0.77	0.39	3	2MASS
HIPASS J1231+20	J123142.4+202853	1332	69	5.1	15.6	0.12	1.20	9	CGCG129-006
HIPASS J1233−00	J123307.9−003159	729	43	3.0	17.28	0.10	3.37	8	UA285=KDG155
HIPASS J1239−04	J123901.9−043352	2412	69	4.8	16.14	0.12	1.85	8	PGC 042298
HIPASS J1239−07	J123944.7−070519	929	53	1.9	16.35	0.10	0.91	10	APMUKS
HIPASS J1242−77	J124206.5−775530	2685	53	3.8	17.7	1.71	1.43	10	
HIPASS J1244−08	J124513.3−082131	2878	84	7.5	15.5	0.15	1.57	8	
HIPASS J1248−46	J124817.8−461304	3414	279	5.8	16.28	0.41	1.96	8	ESO 268-G042
HIPASS J1248−45b	J124904.5−451120	2205	78	2.4	17.0	0.41	1.57	8	
HIPASS J1252−31	J125235.2−315314	3549	132	8.0	15.18	0.37	1.02	8	ESO 442-028
HIPASS J1258−45	J125803.5−454830	2082	70	4.3	16.4	0.39	1.64	8	KKs52
HIPASS J1303−52	J130306.6−521129	1310	64	4.2	17.4	1.58	1.34	10	
HIPASS J1304−02	J130446.5−025217	1271	78	5.6	16.37	0.11	2.70	10	2dFGRS
HIPASS J1305−51	J130504.2−514221	4095	130	4.8	17.2	1.13	1.94	7	PGC 045237
HIPASS J1305−45	J130517.6−453925	3066	53	2.7	16.9	0.39	1.64	8	
HIPASS J1312−15	J131226.3−154752	2953	112	7.8	14.01	0.28	0.37	1	NGC 5010
HIPASS J1314−21	J131409.4−213942	2993	47	2.5	16.1	0.62	0.59	7	ESO 576-G015
HIPASS J1315−22	J131448.2−220450	1399	91	4.7	17.5	0.58	4.17	8	
HIPASS J1316−50	J131601.4−501620	3808	314	18.3	15.4	0.81	1.91	5	ESO 219-43
HIPASS J1320−14	J132013.0−142732	2750	45	4.2	16.5	0.38	1.79	10	pec,[MMB2004]
HIPASS J1321−31	J132108.3−313145	571	39	5.6	17.1	0.27	4.57	9	KK195
HIPASS J1321−31b	J132206.3−312442	2252	195	5.0	16.6	0.27	2.58	8	
HIPASS J1326−09	J132551.0−093219	1803	98	5.3	16.7	0.21	3.16	9	
HIPASS J1328−46	J132738.0−460909	1391	47	4.1	17.6	0.49	4.33	10	
HIDEEP J1328−28	J132823.1−281817	2053			16.94	0.25		10	
HIDEEP J1333−33	J133315.4−334737	2340			16.4	0.25		9	

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J1333−09	J133401.1−090900	2915	149	5.9	16.1	0.17	2.11	8	
HIPASS J1334−12	J133439.7−121950	1511	76	2.2	15.0	0.24	0.27	8	KDG227
HIDEEP J1336−33	J133700.0−332147	591			17.3	0.26		10	
HIDEEP J1337−31	J133658.9−311905	1693			18.8	0.26		5	6dF
HIDEEP J1337−28	J133718.2−283929	2361			15.5	0.25		4	IC 4303
HIPASS J1338−30	J133808.0−305506	1667	135	6.4	16.0	0.24	1.96	9	bluish,2MASS,diffuse tail(?!)
HIDEEP J1338−30	J133845.3−305406	3650			15.93	0.24		4	ESO 444-090
HIPASS J1340−31	J134020.9−314204	1658	96	5.2	16.57	0.24	2.68	8	ESO 445-007
HIPASS J1342−48a	J134200.4−481837	1372	114	7.1	14.4	0.40	0.43	7	ESO 220-032
HIPASS J1342−19	J134246.9−193454	1410	54	2.7	16.42	0.47	0.98	10	ESO 577-27
HIPASS J1345−35	J134455.4−352450	1018	52	2.6	16.01	0.25	0.79	10	ESO 383-70
HIPASS J1348−37	J134833.5−375805	501	39	2.5	18.0	0.33	4.41	10	
HIPASS J1350−37	J135033.3−371831	1077	146	11.0	14.3	0.33	0.64	7	ESO 383-091
HIPASS J1351−47	J135118.9−465825	528	38	3.5	17.5	0.62	2.99	10	
HIPASS J1354−49	J135407.2−492750	1788	103	5.1	17.0	0.83	2.27	8	
HIPASS J1357−46	J135739.2−462618	2850	126	3.7	16.9	0.55	1.94	8	
HIPASS J1400−06	J135958.8−062808	2604	99	3.2	16.18	0.14	1.26	9	APMUKS
HIPASS J1401−05	J140131.8−054933	1564	128	4.3	16.49	0.12	2.29	7	APMUKS
HIPASS J1402−48	J140312.0−483739	2807	142	5.0	16.4	0.81	1.31	7	
HIPASS J1405−51	J140539.9−510445	4092	111	9.7	17.2	1.53	2.73	8	
HIPASS J1407−75	J140729.2−755314	2821	132	7.7	16.8	0.72	3.16	8	
HIPASS J1409−14	J140941.2−144620	2328	106	3.7	16.4	0.35	1.47	7	
HIPASS J1424−34	J142352.1−341529	3118	45	2.4	16.82	0.53	1.19	8	ESO 385-020
HIPASS J1425−39	J142549.0−391603	1174	57	4.5	17.0	0.48	2.75	10	
HIPASS J1428−08	J142826.8−085524	1541	77	5.4	16.9	0.24	3.77	10	KKSG46
HIPASS J1433+01	J143353.5+012912	1827	66	3.2	17.4	0.20	3.66	10	
HIPASS J1435−17	J143525.4−171001	1574	113	9.1	17.2	0.32	7.80	10	KKSG47
HIPASS J1444−17	J144457.4−171453	2826	105	5.5	16.6	0.50	2.29	8	
HIPASS J1452−50	J145256.6−505352	1421	122	5.9	17.5	2.25	1.13	7	ESO 223-03
HIPASS J1546−32	J154626.3−323745	2231	63	5.6	16.53	0.46	2.27	10	ESO 450-12
HIPASS J1548+16	J154902.8+164249	2052	94	5.3	18.0	0.16	10.96	8	

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J1551−70	J155018.5−702854	3328	140	5.6	16.0	0.61	1.21	6	2MASX
HIPASS J1604−75	J160302.5−753932	2912	140	3.1	16.5	0.28	1.45	8	ESO 042-016
HIPASS J1605−04	J160540.8−043420	1611	91	7.2	16.2	1.19	1.10	10	KKSG48
HIPASS J1615−17	J161545.5−175028	2371	197	8.0	16.0	1.51	0.76	8	VLSB,2'x1',cirrus-like
HIPASS J1700−12	J170006.3−120001	1298	77	5.7	17.3	1.23	2.31	8	
HIPASS J1700−07	J170008.1−072324	1558	56	4.9	17.7	2.38	1.00	10	
HIPASS J1711−47	J171137.0−473604	2182	219	20.1	17.6	2.97	2.17	6	
HIPASS J1730−60	J173100.2−601138	1565	46	3.3	17.4	0.37	3.25	8	
HIPASS J1738−57	J173842.9−571525	858	75	4.5	17.5	0.41	4.66	10	
HIPASS J1739−51	J173910.1−510457	3802	146	10.0	17.8	0.71	10.38	8	
HIPASS J1750+21	J175011.5+211556	3222	82	2.2	16.4	0.40	0.84	6	2MASS
HIPASS J1752−59	J175250.9−594057	2736	46	3.3	17.7	0.40	4.17	10	
HIPASS J1758+14	J175839.5+144658	2957	134	4.1	16.8	0.52	2.01	6	
HIPASS J1809−05	J180955.0−055423	2935	250	6.6	18.8	4.94	0.35	7	2MASS
HIPASS J1810−01	J181042.1−010427	2092	165	5.8	18.5	4.87	0.25	7	
HIPASS J1812−74	J181220.0−744957	3216	76	3.8	17.8	0.60	4.37	10	in a star halo
HIPASS J1819+01	J181923.8+011042	2571	114	12.8	18.2	4.35	0.67	6	
HIPASS J1824−01	J182500.2−012833	2875	351	30.5	19.0	6.95	0.30	4	
HIPASS J1832+06	J183214.6+062530	2823	91	7.0	17.7	2.06	1.91	7	
HIPASS J1859−57	J185950.4−570126	3461	42	2.3	16.4	0.32	0.94	6	
HIPASS J1915−57	J191535.4−570559	2991	61	2.5	16.8	0.30	1.50	7	
HIPASS J1919−33	J191932.8−334603	2078	100	3.1	17.0	0.38	2.09	7	
HIPASS J1926−74	J192729.4−740454	2624	43	2.3	17.2	0.34	1.92	9	
HIPASS J1932−55	J193216.5−555429	3413	203	13.0	13.39	0.27	0.35	0	NGC 6799
HIPASS J1937−52	J193738.5−520433	3157	199	21.5	14.5	0.24	1.64	9	IC 4875
HIPASS J1937+09	J193744.4+092125	3148	81	3.3	17.5	1.92	0.86	7	
HIPASS J1951+01	J195157.3+013140	1275	87	3.9	17.7	1.07	2.65	8	
HIPASS J2001−47	J200156.5−471550	3490	90	3.3	15.74	0.18	0.84	8	ESO 284-010
HIPASS J2011−09	J201059.2−095545	2868	86	5.6	17.4	0.39	5.40	8	6dF
HIPASS J2010−11	J201047.6−113829	3204	100	6.9	16.0	0.56	1.57	7	

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
HIPASS J2015+12	J201548.6+124127	1951	52	4.6	18.3	0.87	6.55	10	
HIPASS J2041−16	J204118.7−160424	3452	99	6.4	16.3	0.20	2.68	5	[DDR2005]
HIPASS J2045−53	J204513.7−534353	3255	112	5.8	15.89	0.21	1.64	9	APM
HIPASS J2100−55	J205956.9−553343	2033	188	8.8	13.96	0.21	0.42	2	NGC 6990
HIPASS J2111−50	J211156.8−502111	2790	86	4.1	16.2	0.11	1.69	8	ESO 235-073
HIPASS J2138−54	J213759.5−543130	2924	96	5.7	16.9	0.09	4.57	8	APM, blue,in a group?
HIPASS J2149−35	J215010.9−354228	2458	104	6.5	16.89	0.10	5.11	10	ESO 403-036
HIPASS J2150−23	J215050.5−230848	2342	156	5.2	17.0	0.17	4.25	9	2MASX
HIPASS J2156−49	J215556.9−491438	1865	91	3.9	17.34	0.10	4.66	8	APMUKS
HIPASS J2214−67	J221513.9−672636	3232	80	1.7	15.4	0.14	0.33	4	FAIRALL1005
HIPASS J2217−42	J221654.8−424512	2198	49	2.9	17.0	0.06	2.63	10	AM
HIPASS J2229−79	J223020.5−794131	2531	154	8.3	15.06	0.55	0.80	7	ESO 027-012+companion
HIPASS J2250+00	J225024.4+005241	1692	142	2.8	17.2	0.42	2.19	10	APM
HIPASS J2251−20b	J225148.7−203629	3145	328	14.8	12.63	0.14	0.22	2	NGC 7392
HIPASS J2307−61	J230720.8−614050	3223	48	2.6	17.5	0.09	3.63	10	VLSB
HIPASS J2308+17	J230851.4+171236	1764	142	16.7	15.8	0.56	3.16	8	
HIPASS J2311−42	J231110.9−425051	1381	94	3.4	15.72	0.04	0.96	8	ESO 291-003
HIPASS J2323−50	J232359.4−503127	1579	131	5.1	14.99	0.05	0.73	1	ESO 240-02
HIPASS J2340−20	J234032.6−205639	1397	114	5.4	16.51	0.11	2.96	7	ESO 606-01+companion
HIPASS J2344−07	J234429.7−073547	1832	76	5.7	16.05	0.14	2.00	8	
HIPASS J2358+04	J235807.3+044905	3035	111	4.2	17.9	0.21	7.59	8	
HIPASS J2359+02	J235913.4+024324	2616	148	4.6	16.4	0.12	2.27	6	




Рис. 1: Digitized Sky Survey images of four galaxies identified with HIPASS sources: (a) HIPASS J0727+04, (b) HIPASS J1435-17, (c) HIPASS J2011-09, and (d) HIPASS J2149-35. The size of each field is 6×6 arcmin; north is at the top and east is to the left.

Рис. 2: Distribution of the identified HIPASS sources in hydrogen mass-to-blue luminosity ratio in units of the solar mass and luminosity.

Рис. 3: Hydrogen mass-to-luminosity ratio versus morphological type for 230 HIPASS sources.

Рис. 4: Hydrogen mass-to-luminosity ratio versus observed line FWHM W_{50} for 230 HIPASS sources.

Рис. 5: Distribution of the identified HIPASS sources (circles) in absolute magnitude and hydrogen mass-to-luminosity ratio. The statistical regression line (1) for them is indicated by the solid straight line. Its slope corresponds to the wellknown correlations between the hydrogen mass, radius, and luminosity of disk galaxies. Other nearby galaxies with a high HI abundance are indicated by the triangles. The dashed line indicates the minimum stellar mass fraction required to maintain the gravitational stability of a galaxy according to Warren et al. (2007). The cross marks a typical error of our apparent magnitude estimate for the galaxies.

This figure "Karachentsev_fig2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0902.0256v1>

This figure "Karachentsev_fig3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0902.0256v1>

This figure "Karachentsev_fig4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0902.0256v1>

This figure "Karachentsev_fig5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0902.0256v1>